

Slow-release nitrogen fertilizers effects on plant growth and yield of sorghum

Mona E. El-Azab

Soil and water Dept., National Research Center, Dokki, Egypt

Abstract : The aim of this investigation was to study the efficiency of conventional nitrogen fertilizers compared with slow-release fertilizers as urea, ureaform and sulfur- coated urea (SCU) at different doses on sorghum plants. Field experiment was conducted at Al-Sharkia Governorate, Egypt in a private farm through a project of soil and water use Dept. of the National Research Center. This experiment design with three replicates. Sorghum seeds were sown on the 15th of December, 2015. Plants were fertilized by Seven treatments of nitrogen fertilizer from control (no fertilization), (urea 46%N), (ureaform 38%N) and (SCU 32%N) respectively, with two levels of nitrogen as 100 and 200 Kgfed⁻¹. Results showed that the use of slow-release nitrogen fertilizer gave the highest values of vegetative growth as plant height, number of leaves/plant, number of internodes/plant and dry weight and increase the yield and its components beside increase the concentrations of N, P and K and its uptake at all growth stages. The use of sulfur coated urea more effective than ureaform of the parameters under investigated.

Keywords : urea, slow-release fertilizers, nitrogen efficiency, sorghum.

Introduction

Sorghum (*Sorghum bicolor* L.) plant a cereal crops is very important for economically after corn, wheat and rice¹. This plant can be grown under hard situations such as soil moisture, temperature and it responses to irrigation very good so it can grown in different zones of the world. For plant growth of sorghum, nitrogen is very necessary to improve the growth and yield¹¹. Nitrogen is the major factor for increased or reduced yields³³. Nitrogen is important as macronutrients for increase plant growth. Its main effect are use as the component of many important organic compounds and collaborate in many essential metabolic processes in plants such as building amino acids, proteins, carriers, enzymes, regulators, nucleic acids, pigments, alkaloids and many other metabolites^{20,29}. More studies effects the nitrogen fertilization on plant growth such as¹² obtained that the use of nitrogen fertilizers increase the plant growth and yield. But the large use of nitrogen fertilizers increase the production costs and it causes the pollution of environment^{25,19}.

Nitrogen use efficiency is significance in crop production system because its effect on farmer economic outcomes and environmental influence. Also, may be decrease the crop production resulted to many factors as losses of soil nitrogen by volatilization, leaching and denitrification. Conducted numerous research to increase sorghum yield by increasing the efficiency of fertilizers use and increasing of nutrient absorption by plants through use the controlled- release fertilizers which can be reduce the nutrients loss by leaching and runoff¹⁷. Slow-release nitrogen fertilizers is very important to save nitrogen from loss by leaching from plant, it can be increase the level of nitrate (NO₃) in ground and surface water¹⁸. Recently, slow-release nitrogen fertilizer has

been suggest as to increase N fertilizer for sandy soil that loss N leaching and increase N use efficiency ²¹. Slow-release fertilizers are new technique to increase the nutrient availability to plant as well as loss the release of nutrients to soil compared to another conventional fertilizers¹⁷. ⁸ showed that the use of slow-release nitrogen fertilizer improved the plant growth and gave the highest yield of sorghum plants compared to other nitrogen sources. ²⁶ suggested that the use of slow-release N fertilizers can be decrease N leaching and increase plant growth which increase nitrogen concentration in plant. ³³ showed that after application of slow-release fertilizer up to 30% of N applied that can be leached compared to more 88% N leaching after application the soluble ammonium nitrate in sandy soils. In general, the major advantages for coating fertilizers are that they make an acidic environment from acidic materials used and inhibit ureolytic microorganism's activities which in effect slows down the release of NH_4^+ into the soil and indirectly decrease N loss ⁵.

Slow-release fertilizers showed that the release of the nutrient in a lower quantity than other fertilizers. However, slow-release fertilizers are different in rate, form and period of release. It will be strongly affected with storage, transportation and distribution in the field, or by soil conditions such as moisture content, wetting and drying, thawing and freezing, and biological activity ^{5,27,23}.

The main role of slow-release nitrogen fertilizer that improving the efficiency of fertilizers and decrease the loss of nitrogen. This result obtained with ¹⁴ showed that the use of slow-release fertilizers improve the plant growth and the coating materials decrease or delaying the release of nitrogen from fertilizers granules to soil. ¹⁶ who obtained that the application of slow-release nitrogen fertilizer increasing yield of sorghum. ^{24,36} also indicated that the application of slow-release with urea increased wheat yields from 18.3 – 27.8 % and rice yields from 27.5 – 50.4 % than application of urea only. ²¹ found that the slow-release N can decrease N loss by leaching, increase plant growth and increase N concentration compared with ammonium nitrate. Slow-acting fertilizers are characterized by a delay between the time of application and nutrient availability, as well as a lower release of nutrients into soil compared to conventional fertilizers¹⁷.

There is some information of use slow-release fertilizers on sorghum. Therefore the present study was conducted to evaluate the effect of nitrogen fertilizers with slow-release fertilizers on yield and some nutrients uptake of sorghum plants.

Materials and Methods

A field experiment was executed in Sharkia Governorate, Egypt, to effects the efficiency of nitrogen fertilizers with slow-release fertilizers with two levels on nutrient concentration and yield of sorghum plant (*Sorghum bicolor* L.). Sorghum seeds were sown on the 15th of December, 2015. Superphosphate (15.5% P O) was added at rate of 150 Kg/fed before sowing. Plants were fertilized with Seven treatments of nitrogen fertilizer as control, urea (46%N), ureaform (36%N), SCU (32%N) respectively, with two levels of nitrogen as 100 and 200 Kgfed⁻¹. The fertilizers were added before irrigation which applied at 15, 45 and 60 days from sowing. Treatments were as follows:

1. Control
2. Urea 100Kgfed⁻¹
3. Urea 200Kgfed⁻¹
4. Ureaform100 Kgfed⁻¹
5. Ureaform200 Kgfed⁻¹
6. SCU100 Kgfed⁻¹
7. SCU200 Kgfed⁻¹

The physical – chemical properties of the soil is presented in Table (1). Chemical formula and nitrogen percent of the applied slow-release fertilizers and conventional fertilizers is presented in Table (2). Treatments were arranged in a complete randomized block design with 3 replicates. Plant samples were taken at 60, 90 and 120 days after planting. Plant height (cm), number of leaves/plant, number of internodes, dry weight of plant, grain weight, shoot and grain yields as well as 100-grain weight were recorded and grains were also subjected to chemical analysis. Grain weight kg/plot were measured. Grain yield per feddan was obtained. Total of nitrogen, Phosphorus and potassium were determined.

Methods of Analysis

Mechanical analysis was determined according to the international Pipette method and calcium carbonate content of the soil was determined volumetrically using Calcimeter as described by ²².

The electrical conductivity (EC) of soil water extract was determined by using the bridge, ¹³.

Calcium carbonate content of the oil was determined volumetrically using Collins calcimeter as described by ²².

Soil pH was measured using a glass electrode pH meter in a 1: 2.5 soil water suspension ⁶.

Organic matter was determined by Walkley and Black's method as described by ¹³.

Soluble cations and anions were determined in (1:5) soil water extract according to ⁴.

Sodium and potassium were determined by using flame photometer as described by ⁶.

Calcium and magnesium were determined following the versenate method ¹³.

Total nitrogen in soil was determined using microKjeldahl method ¹³.

Total potassium in soil was determined by flame photometer according ¹³.

Total phosphorus in soil was determined colourmetrically using ascorbic acid method ³².

The data obtained was subjected to analysis variance procedure using Duncan's Multiple Range Test was adopted for the means comparison among treatments showing significant difference. Effect of N and P fertilizer was partitioned into linear and quadratic components and regressions were calculated for effects significant at 0.05 level of probability ⁷.

Table (1): Physical and chemical properties of the studied soil

Soil characteristics	Soil content
Mechanical analysis:	
Fine sand%	12.36
Coarse sand%	66.42
Silt%	6.80
Clay%	14.42
Textural	Sandy
Chemical analysis:	
Organic matter%	1.04
pH	7.31
EC (dS m ⁻¹)	0.12
CaCO ₃ g kg ⁻¹	2.25
Soluble ions (mmol⁻¹)	
Ca ⁺⁺	0.34
Mg ⁺⁺	0.56
Na ⁺	0.66
K ⁺	0.12
CO ⁻³	-
HCO ⁻³	0.37
Cl ⁻	0.48
SO ⁻⁴	0.29
Available-N (g kg ⁻¹)	2.47
Available-P (g kg ⁻¹)	1.22
Available-K (g kg ⁻¹)	0.63

*Soil- water suspension 1:2.5 ** Soil water extract 1:5

Table (2): Chemical formula and nitrogen percent of the applied slow-release fertilizers and conventional fertilizers.

Material	Formula	N%
Urea	$\text{O}=\text{C}(\text{NH}_2)_2$	46
Ureaform	$\text{HO}-\text{CH}_2-\text{CO}-\text{NH}-\text{CH}_2-\text{NH}-\text{C}=\text{O}-\text{H}$	36.2
Sulfur-coated urea "SCU"	$\text{O}=\text{C}(\text{NH}_2)_2+\text{S}$	32

Results and Discussion

Vegetative growth:

Results show that the effect of nitrogen fertilizers of sorghum plants with slow-release fertilizers at different concentrations on vegetative growth. Concerning the effect of nitrogen fertilizers with slow-release fertilizers on vegetative growth at 60 and 90 day after sowing, data presented in Table (3), showed that the slow-release fertilizers was generally more effective than the control plants and urea. These results may be due to the effect of coating materials on decrease of nitrogen loss from fertilizers. These results were in agreement with the results obtained by ¹⁷ and ⁸ who revealed that the application of slow-release nitrogen fertilizer increased the growth of sorghum plants compared to other nitrogen sources. Data obtained that the application of Slow- release N fertilizer as ureaform and sulfur coated urea gave the highest values of plant height, number of leaves/plant, number of internodes/plant and dry weight of plant compared with urea and control. This result may be attributed to the effect of coating materials on delaying the release of nitrogen from fertilizer granules as indicated by ¹⁴ and ³¹ who obtained that coating materials can decrease nitrogen loss through intermittent leaching by stalling the release of nitrogen. Data presented in the same Table showed that sorghum grown in soil fertilized with slow-release N fertilizer improved the plant growth at 60 and 90 day after sowing, than the other sources of N fertilizer as urea. Result may be due to the solubility of urea was greater than ureaform and sulfur coated urea so the application of urea increased the loss of nitrogen compared with slow-release fertilizer. These results are agreement with ^{31,17,26}.

The maximum mean values of plant height, number of leaves/plant, number of internodes/plant and dry weight of plant at 60 and 90 day after sowing, were found with foliar slow-release fertilizer with 200Kgfd^{-1} . On the other hand, the minimum values of the vegetative growth were occurred with urea fertilizer with 100Kgfd^{-1} . These results may be due to the effect of slow-release fertilizers in plant physiology which it increase the nitrogen uptake from plant that nitrogen plays a important role in several vital processes in plant such as photosynthesis consequently affecting plant growth which was effect on vegetative growth plant. Improving effect of slow-release fertilizers on vegetative growth might be attributed to their positive role on nitrogen loss and it was increase the rate of photosynthesis, biosynthesis of proteins and carbohydrate assimilation. This result in coincidence with the findings of ²⁶. Data in the same Table, illustrate that slow-release fertilizers treatments with 200Kgfd^{-1} increased significantly the dry weight compared with urea and control. The positive effect of slow-release fertilizers was greater than urea. These result may be due to the solubility of urea was greater than urea form and sulfur coated urea. These results are agreement with ³¹. Data in the same Table showed that the maximum mean values of vegetative growth at 60 and 90 day after sowing, were obtained with sulfur coated urea at 200Kgfd^{-1} than urea form. This result may be attributed to the low activity index of urea form as recorded by ³.

Table (3): Effect of different sources of nitrogen fertilizers on vegetative growth of sorghum plant.

Treatments		plant height (cm)		number of leaves/plant		number of internodes/plant		dry weight (g)	
fertilizers	Rate of N Kg/fed	60	90	60	90	60	90	60	90
Control	0	98.24	110.28	4.33	4.00	5.00	5.00	94.12	115.36
Urea	100	104.24	129.34	5.67	5.33	6.00	6.00	127.45	133.54
	200	114.36	138.55	6.00	5.50	6.00	6.00	128.23	142.58
Ureaform	100	126.22	135.39	7.00	6.50	7.00	7.00	145.22	154.23
	200	131.25	144.21	7.50	7.00	7.00	7.00	151.14	156.91
SCU	100	132.69	146.11	7.31	7.00	8.00	8.00	156.35	163.41
	200	134.41	148.57	7.66	7.50	8.00	8.00	158.14	164.54
LSD at 0.05		1.26	1.44	0.11	0.24	0.25	0.16	1.48	1.23

Yield and its components:

Data in Table (4) obtained that the grain weight, shoot and grain yields as well as 100-grain weight as affected by nitrogen fertilizer with slow-release fertilizers under experimental conditions. The maximum values of this characteristics were produced from slow-release fertilizers application. While, the minimum values of aforementioned attribute were obtained with control. These results are in harmony with those obtained by⁸ who suggested that there are several potential benefits of providing nitrogen to slow-release fertilizer. These include: reduced nitrogen losses through denitrification and leaching compared with nitrogen fertilizer application as urea. The superior of slow-release than urea can be attributed to the slow-release of N to meet plants requirement, where it has a low dissolution rate than the others which reduces nitrogen loss from soil profile and gives a chance for more nitrogen uptake by plant root. Similar finding were reported by¹⁶ who found that slow-release nitrogen fertilizer gave the highest yield of sorghum. ²⁴also showed that slow-release urea increased wheat yields by 18.3 – 27.8 % and rice yields by 27.5 – 50.4 % as compared with common urea. ²¹ showed that amending sandy soils with slow-release N can reduced leaching, increase plant growth and increase N concentration compared with sweet corn grown in soil amended with ammonium nitrate. ²⁸ told that, slow-release N fertilizer overcame soluble N fertilizer.

Sorghum grown in soil fertilized with slow-release N fertilizer as ureaform and sulfur coated urea gave significantly more yield and its components. These result due to effect of slow-release fertilizers on nitrogen loss which increase the plant growth and effect on photosynthesis rate. These results are agreement with ^{31,17,26}. Although sorghum uses nitrogen more efficiently than most plants with the C3 cycle, nitrogen is the main factor responsible for reduced yield³⁴.

Data showed that all slow-release fertilizers treatments with 200Kgfed⁻¹ greatly improved the yield and its components of sorghum plants compared with control, grain weight, shoot and grain yields as well as 100-grain weight were significantly improved with increasing of slow-release fertilizers with rate from 100Kgfed⁻¹ up to 200Kgfed⁻¹. These results may be due to the effect of these slow-release fertilizers on nitrogen loss so it effect on plant physiology such as photosynthesis consequently affecting plant growth, important for ¹⁴C fixed in the primary photosynthetic process. These increases of the grain weight, shoot and grain yields and 100-grain weight may be due to the amount of metabolites synthesized by plants as a result of increasing nitrogen levels. This may be attributed to the favorable effect of nitrogen fertilizer levels on the metabolic processes and physiological activates of meristematic tissues, which are responsible for cell division and elongation in addition to formation of plant organs this lead to more vigorous growth and consequently accumulation of more photosynthesis assimilates. Similar results were reported by^{10,9}.

The obtained results are in harmony with those of² and ¹⁵ where they found that yield and its components increased markedly by slow-release fertilizers with 200Kgfed⁻¹ compared with the untreated plants.

Although sorghum uses nitrogen more efficiently than most plants with the C_3 cycle, nitrogen is the main factor responsible for reduced yields³⁴. So, the uses of slow-release fertilizers decrease of nitrogen loss and increase the nitrogen uptake from plant.

It is clear from Table (4) that slow-release N fertilizer as sulfur coated urea at 200kgfed^{-1} was the best treatment in all characters under study especially grain yield and straw yield/plant compared with urea. This result may be attributed to the loss nitrogen from urea. The positive effect of slow-release fertilizers was greater than urea. These result may be due to the solubility of urea was greater than ureaform and sulfur coated urea. This result are agreement with³¹. Data in the same Table showed that the maximum mean values of yield were obtained with sulfur coated urea at 200Kgfed^{-1} compared with ureaform. This result may be attributed to the low activity index of ureaform as recorded by³. On the other hand, the minimum mean values of grain yield were obtained with control. Similar finding were obtained by^{28,8}. Slow-release N fertilizer may help positively in plant growth, which was increased of N, P and K content in plant because it saved the nutrients from loss. Similar results were reported by^{10,9}.

Table (4): Effect of different sources of nitrogen fertilizers on yield and its components of sorghum plant at harvest.

Treatments		grain weight/ plant(g)	shoot yield /plant(g)	grain yields (ton/fed)	shoot yield (ton/fed)	100-grain (g)
fertilizers	Rate of N Kg/fed					
Control	0	24.11	114.24	0.87	4.24	22.47
Urea	100	34.58	186.24	1.46	8.74	25.14
	200	38.22	195.23	1.51	8.94	26.32
Ureaform	100	41.26	214.56	2.47	9.35	28.59
	200	44.59	232..14	2.63	9.47	28.45
SCU	100	48.26	241.58	2.57	9.66	30.18
	200	51.23	249.36	2.78	9.71	30.53
LSD at 0.05		1.22	1.24	0.45	0.33	1.29

Nutrient concentration:

Results showed that the application of slow-release fertilizer with different levels on nitrogen, phosphorus and potassium concentration (%) and its uptake (g plant^{-1}) in sorghum grains. The results in Table (5) show that application of slow-release fertilizer with 200Kgfed^{-1} significantly increased nutrient concentration of sorghum plant compared to control treatment or urea application. This result may be due to effect of slow-release fertilizers on nutrients loss which decrease the loss by leaching so effect on translocation rate of photosynthesis. These results are in agreement with those obtained by³⁰ who showed that the application of slow-release fertilizers increased the N, P and K concentration in plant.

The results indicate that the highest concentration of nitrogen and it is uptake with slow-release fertilizers as urea form and sulfur coated urea at 200K gfed^{-1} compared with the application of urea, but the lowest concentration of nitrogen at control. Also, the results show that the highest concentration of phosphorus and potassium and it is uptake absorbed with ureaform and sulfur coated urea at 200K gfed^{-1} compared with treatment urea. And the lowest concentration of phosphorus and potassium and it is uptake absorbed with control. These results may be attributed to effect of slow-release fertilizers to reduce N leaching and increases the yield of sorghum in such sandy soil. Also the effect of slow-release fertilizers is positive effect from nutrients availability for plant. This result obtained by²¹ showed that amending sandy soils with slow-release N can reduced leaching, increase plant growth and increase N,P and K concentration compared with sweet corn grown in soil amended with ammonium nitrate. ²⁸ showed that, the slow-release fertilizer increase the nutrients availability to plant which decrease the nutrient loss by leaching. It might be due to participation of slow-

release fertilizers with 200Kg fed^{-1} which reduce the nitrogen loss so, its plays important roles in plant growth and production, including chlorophyll synthesis, protein synthesis and chloroplast development.

From data in the same Table, the results indicate that the highest concentration of nitrogen, phosphorus and potassium and its uptake in sorghum grains were obtained with sulfur coated urea than urea form. This result may be attributed to the low activity index of urea form as recorded by³.

Nitrogen Efficiency Ratio (NER) and Nitrogen Recovery (NR)

Data in the Table (6) showed that the effect of nitrogen fertilizers and slow-release fertilizers on nitrogen efficiency ratio (NER) and nitrogen recovery (NR). The results of Table obtained that there are significantly differences between nitrogen efficiency ratio and nitrogen recovery in sorghum grains. Also, in the same Table, the highest mean value of (NER) obtained with slow-release fertilizers as ureaform and sulfur coated urea compared with urea and control. The lowest mean value of NER observed with control. Data of nitrogen recovery (NR) which presented in Table (6) show that the highest (NR) values has been obtained from slow-release fertilizers. While, the lowest values of (NR) has been obtained with control and urea. This result may be due to effect of slow-release fertilizers on nitrogen loss which decrease the loss by leaching. These results are in agreement with those obtained by³⁰. Regarding with the effect of slow-release fertilizers rates, data in the same Table noticed that there are significantly differences between nitrogen efficiency ratio and nitrogen recovery as affected by slow-release fertilizers rates. Also, the highest mean value of NER was observed with sulfur coated urea at 200Kg fed^{-1} . Meanwhile, the lowest mean value of (NER) obtained with urea fertilizer at 100Kg fed^{-1} . These results are in harmony with those obtained by³⁰. Also, the highest mean values of NR was observed with sulfur coated urea at 200Kg . And the lowest values of NR was obtained with urea at 100Kg fed^{-1} . This result may be attributed to the loss nitrogen from urea. The positive effect of slow-release fertilizers was greater than urea. These result may be due to the solubility of urea was greater than urea form and sulfur coated urea. This result are agreement with³¹. Similar results were reported by^{10,9}.

Data in the same Table showed that the maximum mean values of (NER) and (NR) were obtained with sulfur coated urea at 200Kg fed^{-1} compared with ureaform. This result may be attributed to the low activity index of ureaform as recorded by³. On the other hand, the minimum mean values of (NER) and (NR) were obtained with urea. Similar finding were obtained by^{28,8}.

Table (5): Effect of different sources of nitrogen fertilizers on N, P and K concentration and its uptake of sorghum grains.

Treatments		%			Uptake (g plant^{-1})			NER	NR
fertilizers	Rate of Kg/fed	N	P	K	N	P	K		
Control	0	1.46	0.12	0.34	0.35	0.03	0.08	8.44	14.23
Urea	100	1.68	0.23	1.33	0.58	0.08	0.46	20.06	18.67
	200	1.87	0.25	1.42	0.71	0.09	0.54	27.14	19.47
Ureaform	100	2.14	0.34	1.75	0.88	0.14	0.72	36.31	21.65
	200	2.21	0.41	1.88	0.98	0.18	0.83	43.70	22.14
SCU	100	2.32	0.42	1.89	1.12	0.20	0.91	54.05	22.35
	200	2.54	0.45	1.91	1.30	0.23	0.97	66.59	24.57
LSD at 0.05		0.15	--	0.24	2.13	0.39	1.58	1.22	0.12

Conclusions:

There were significant differences between nitrogen fertilizers and slow-release fertilizers. For sandy soil, the use of slow-release nitrogen fertilizers was very important to reduces N leaching and increases the yield of sorghum. Also the use of slow-release N is more effective than another nitrogen fertilizers generally resulted in increased the growth and nutrients uptake by sorghum plants grown in sandy soil. Using slow-

release fertilizers reduces production costs, eliminates the need for multiple applications of soluble nitrogen fertilizers. Also slow release nitrogen fertilizers were able to increase nitrogen use efficiency by reduce nitrogen by leaching and volatilization from soils.

Acknowledgment

Many thanks to the staff members of soil and water use Department National Research Center for great help During This Study.

References

1. Agricultural Economic Year Book. Ministry of Agric. A. R. C. 1995.
2. Amal, G. A., N. Zaki, N and Hassanein, M. Response of grain sorghum to different nitrogen sources. Research Journal of Agriculture and Biological Sciences, 2007, 3(6): 1002- 1008.
3. Awad, E. Effect of vetera huudrogel and nitrogen fertilizers on wheat. Zagazig. J. Agric. Res. 1990, 17: (48).
4. Black, C.A. Methods of Soil Analysis. Part I- Physical and mineralogical properties. A. S. A. Madison Wise., USA.1965.
5. Cheftetz, B., P.H. Hatcher, P. H., Hadar Y and Chen, Y. Chemical and biological characterization of organic matter during composting of municipal solid waste. J. Environ. Qual., 1996, 25: 776-785.
6. Cottenie, A., Verloo, M., Velghe, G. and Kiekens, L. Biological and Analytical aspects of soil pollution. Laboratory of Analytical and Agrochemistry, State University, Ghent-Belgium. 1982.
7. Duncan, D.B. Multiple Range and Multiple F. Test. Biometrics. 1965, 11, 1-42.
8. El-Kramany, M. F. Effect of organic manure and slow-release N-fertilizers on the productivity of wheat (*Triticum aestivum* L.) in sandy soil. Acta. Agronomica Hungarica, 2001, 49: 379-385.
9. El-Bana, A.Y.A. and Gomaa, M. A. Effect of N and K fertilization on maize grown in different populations under newly reclaimed sandy soil Zagazig J. Agric. Res., 2000, 27(5): 1179-1190.
10. El-Naggar, M.A. and Amer, E. A. The effect of nitrogen fertilizer on some maize cultivars in relation to the yield and the infestation by *Ostrina nubilalis*. Minufiya J. Agric. Res., 1999, 24(3): 937-943.
11. Eweis, E.G., Mohamed, K. A and Rayan, A. Interaction effect of number of irrigations and nitrogen rates on grain sorghum. Egypt. J. Appl. Sci. 1998, 13(7): 396-403.
12. Fageria N. K and Moreira A. The role of mineral nutrition on root growth of crop plants. Adv. Agron. 2011, vol. 110: 251-331.
13. Jackson, M. L. Soil Chemical Analysis. Printic Hall Englewood Cliffs, New Jersey.1958.
14. Jarrell, W.M., Pettygrove, G.S. and Boersma, L. Characterization of the thickness and uniformity of the coatings of sulphur coated urea. Soil Sci. Soc. Am. J. 1979, 43, 602-605.
15. Kaneta, Y., Awasaki, H and Murai, Y. The non-tillage riceculture by single application of fertilizer in a nursery box with controlled-release fertilizer. Japanese. Nippon Dojo Hiriyogaku Zasshi, 1994, 65(4), 385-91.
16. Kolhe, S.S and Mittra, B. N. Evaluation of slow-release nitrogen fertilizers in rice-wheat cropping system. Indian J. of Agronomy, 1989, 34: 137-138.
17. Korzeniowska, J. Progress in research on slow-release fertilizers. Stud. Rap. 2009, 18: 9-26.
18. Lichtenberg, E and Shapiro, L. S. Agriculture and nitrate concentrations in Maryland community water system wells. J. Envir. Qual., 1997, 26: 145-153.
19. Marsalis, M., Angadi, S and Contreras-Govea, F. Dry matter yield and nutritive value of corn, forage sorghum, and BMR forage sorghum at different plant populations and nitrogen rates. Field Crop. Res., 2010, 116: 52-57.
20. Marschner, H. Mineral Nutrition of Higher Plants. Second edition, Academic Press, Harcourt Brace & Company, Publishers.1999.
21. Perrin, T.S., Boettinger, J., Drost, D and Norton, J. Decreasing nitrogen leaching from sandy soil with ammonium-loaded clinoptilolite. J.Environ. Qual, 1998, 27: 656- 663.
22. Piper, C. S. Soil Land Plant Analysis Inter. Science Publisher inc., New Yourk.1950.
23. Raban, S., Zeidel, E and Shaviv, A. Release Mechanisms Controlled Release Fertilizers in Practical Use. Dahlia Greidinger Sym. on Fertilization and The Environment, 1997, 287-295.

24. Ragasits, J.B. and Berecz, K. Effect of slow-release N-fertilizers and yield and baking quality of winter wheat. *Fertilizers and Environment*, 1996, 237- 240.
25. Sainju, U., Whitehead, W., Singh, B., Wang, S. and Tillage, S. cover crops, and nitrogen fertilization effects on soil nitrogen and sorghum yields. *Europ. J. Agron.*, 2006, 25: 372-382.
26. Scott Perin, T., T., Danial, L.B., and Jeanette, M. Ammonium-loaded clinoptilolite : A slow-release nitrogen fertilizer for sweet corn. *J. Plant Nutr.*, 1998, 21:515-530.
27. Shaviv, A. Plant response and environmental aspects as affected by rate and pattern of nitrogen release from controlled release N fertilizers. In *Progress in Nitrogen Cycling Studies. Fertilizers and Environment*, 1996, 285-291.
28. Sonbol, H.A., Taha, A., El-Sirafy, Z and El-Naggar, E. Efficiency use of bio and chemical nitrogen fertilizers on wheat. 8th Conf. Agric. Dev. Res. Fac. Agric. Ain Shams Univ. Cairo. 2002, November, 20-22.
29. Srivastava, H. S. and Singh, R. *Nitrogen Nutrition and Plant Growth*. Oxford & IBH Publishing Co. PVT. LTD, 1999.
30. Torbatinejad, N.M., Chajchee, M and Sharifi, S. The effect of different levels of nitrogen on silage quality of three sorghum cultivars. *Agricultural Sciences and Technology*. 2001, 15(2): 75-82.
31. Vallejo, A., Cartagena, M. C., Rodriguez, D. and Diez, J. Nitrogen availability of soluble and slow-release nitrogen fertilizer as assessed by electroultra filtration. *Fert. Res.* 1993, 34: 121-126.
32. Watanabe, F.S. and Olsen, S.R. Acid method for determining phosphorus in water and NaHCO₃ extracts from soil. *Soil. Soc. Am. Proc.* 1965, 29:677-678.
33. Wang, F. L. and A. K. Alva (1996). Leaching of nitrogen from slow-release urea source in sandy soils. *Soil Sci. Soc. Am. J.* 60:1454-1458.
34. Young K.J. and Long S.P. Crop ecosystem responses to climatic change: maize and sorghum. In: *Climate change and global crop productivity*. Reddy K.R., Hodges H.F. CABI Publishing, Wallingford. 2000, 107-131.
35. Zeidan, M.S. and El Karamany, M. Effect of organic manure and slow-release N-fertilizers on the productivity of wheat (*Triticum aestivum* L.) in sandy soil. *Acta Agronomica Hungarica*, 2001, 49(4): 379-385.
36. Zhao, G., Liu, Y., Tian, Y., Sun, Y. and Cao, Y. Preparation and properties of macromolecular slow-release fertilizer containing nitrogen, phosphorus and potassium. *J. Polym. Res.*, 2010, 17:119-125.
