

Effect of *Clerodendrum phlomidis* on adjuvant induced arthritis in rats - A radiographic densitometric analysis

D.Kilimozhi¹, V. Parthasarathy^{1*} and N.Amuthavalli²

¹Department of Pharmacy, Annamalai University, Annamalai Nagar-608002, Tamil Nadu, India

²Department of Radiology, Rajah Muthiah Medical College and Hospital, Annamalai University, Annamalai Nagar-608002, Tamil Nadu, India

*Corres author: vapartha123@gmail.com

Phone: +91-9443512724, Fax: +91-4144-238080

ABSTRACT : In the present study, the anti-arthritic effect of oral administration of ethanolic extract of *Clerodendrum phlomidis* on Freund's adjuvant induced arthritis has been studied in Wistar albino rats. The loss of body weight during the arthritic condition was corrected on treatment with ethanolic extract of *Clerodendrum phlomidis* at 250 and 500 mg.kg⁻¹body weights. The swelling of the paws during the secondary lesions was also markedly reduced on treatment with ethanolic extract of *Clerodendrum phlomidis* and this result was confirmed using radiographic analysis and the changes in the density of Hind Limb Bone Mass (HLBM) was measured using photodensitometer and aluminium step wedge. The HLBM was significantly reduced on treatment with ethanolic extract (250 and 500 mg.kg⁻¹body weight) of *Clerodendrum phlomidis* and standard drug Indomethacin (10 mg.kg⁻¹). From the result we observed that the ethanolic extract of *Clerodendrum phlomidis* possess potent anti-arthritic activity.

KEYWORDS - *Clerodendrum phlomidis*, Anti-arthritic, Freund's complete adjuvant, Photodensitometer, Aluminium step wedge.

1. Introduction

Arthritic affects 0.5-1% of the world population with more women being affected than men [1]. The immune system is a well-organized and well-regulated. The deregulation of the immune system may lead to the development of autoimmune diseases such as Rheumatoid arthritis (RA), is proto-type of such groups of illness with chronic, systemic disorders with destructive inflammatory polyarticular joint potentially resulting in progressive destruction of articular and periarticular structure [2]. Persistent inflammation produces swollen joints with severe synovitis, decreased nociceptive threshold [3, 4] and massive subsynovial infiltration of mononuclear cells, which is along with angiogenesis leads to pannus formation [5]. Expansion of the pannus induces bone erosion and cartilage thinning, leading to the loss of joint function [6, 7]. This result in a high degree of morbidity and disturbed daily life of the patient. Corticosteroids have not been able to fully control the incidence because of its limitations and risk of side effects. Many patients and practitioners are seeking alternative approach to provide an effective cure in the treatment of arthritis and to overcome the serious draw backs such as gastro

intestinal bleeding [8] on treatment with Corticosteroids. Hence there is an urgent need to find safer drug for the management of rheumatoid arthritis.

Clerodendrum phlomidis (L.) Gamble (verbinaceae) is found in some part of south India and it is a fast growing tree in the footing of rivers and channel banks [9-11]. The height of the tree is 6-9m height, tolerably smooth and ash colored. In the Indian system of medicine the leaves of plant is used for rheumatism, anti-microbial [12] and flatulence [13]. Until now no work has been carried out to assess anti-arthritic activity of *Clerodendrum phlomidis*.

Hence the present investigation was undertaken to study the anti-arthritic activity of the ethanolic extract of *Clerodendrum phlomidis* due to the fact that any botanical was traditionally used for wound healing, fever, infection, pain, edema or rheumatic disorders is taken as an indicator that the plant should be tested for its anti-arthritic properties [14].

2. Materials and methods

2.1 Plant Material

Taxonomic identification of the plant was made from Rapinat Herbarium, St. Joseph's college of arts and

sciences, Trichy, Tamilnadu, India. Whole fresh plant leaves of *Clerodendrum phlomidis* was collected from Jeyankondam, Perambalur (Dist), Tamilnadu, India. The leaves were dried under shade, segregated, pulverized by a mechanical grinder and passed through 40 mesh sieves.

2.2 Preparation of extracts

The powdered leaves (1000g) were successively extracted with ethanol (70-80°C) for 24 hrs by continuous hot percolation method using soxhlet apparatus. The fraction was separated from the solvent by distillation under reduced pressure to yield a solid mass (9.6% w/w), stored in refrigerator and used for further studies.

2.3 Animals

The animals for the present study procured after animal ethical clearance from the Institutional Animal Ethical Committee (IAEC) in Annamalai University, Annamalai nagar, Tamilnadu, India. The animal experiments were carried out according to Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) rules. Inbred Wister rats (150-200g) were used for testing anti arthritic activity. The animals were housed at central animal house (Rajah Muthiah Medical College and Hospital, Annamalai University, Tamilnadu, India) under standard conditions of temperature (23±1°C), relative humidity (55±1%) and 12hrs light and dark cycles. The animals were and fed with standard pellet diet and tap water *ad libitum*.

2.4 Drugs and chemicals

All the drugs used in this study were of pharmaceutical grade. Freund's adjuvant was supplied by Genex Pharma, Indomethacin is a gift sample from Cadila Pharmaceuticals, Ahamedabad, India.

2.5 Acute toxicity studies

Acute toxicity studies were performed [15] according to OECD-423 guidelines (acute toxic class method). Albino mice (n=3) of either sex selected by random sampling technique were employed in this study. The animals were fasted for 4hrs with free access to water only. The plant extract of *Clerodendrum phlomidis* was administered orally with an initial dose of 1000 mg.kg⁻¹ body weight. The mortality was observed for three days. If mortality was observed in 2/3 or 3/3 of animals, then the dose administered was considered as a toxic dose. However, if the mortality was observed only one mouse out of three animals then the same dose was repeated again to confirm the toxic effect. If mortality was not observed, the procedure was then repeated with higher dose such as 50,300 and 2000 mg.kg⁻¹.

2.6 Evaluating of paw volume and body weight changes in Freund's induced arthritic animal.

Freund's adjuvant induced arthritis model [16] was used to assess the anti-arthritic activity of the ethanolic extract of *Clerodendrum phlomidis* in Wister rats. Animals were randomly divided into five groups of six animals each (n=6). Group I animals received normal saline (5 mg.kg⁻¹) served as control, Group II animals received Indomethacin (10 mg.kg⁻¹ p.o.) served as reference standard, Group III animals received (0.01 ml Freund's adjuvant) served as an arthritic control and Group IV and

V animals received the crude ethanolic extract of *Clerodendrum phlomidis* (250 and 500 mg.kg⁻¹). The paw volume is a indicator of arthritic condition. To assess the anti-inflammatory and anti-arthritic activity of *Clerodendrum phlomidis*, the extracts were given to the animal 30 minutes before the administration of Freund's adjuvant and continued till 28th day. Paw volume was measured on 4th, 8th, 12th, 16th, 20th, 24th and 28th day by using plethysmometer and changes in body weight also measured.

2.7 Evaluation of radiological changes in the hind limb paw

Freund's adjuvant induced Arthritis model [16] was used to assess the anti-arthritic activity in Wister rats. Animals were randomly divided into five groups of six animals each (n=6). Group I animals received normal saline (5 mg.kg⁻¹) served as control, Group II animals received Indomethacin (10 mg.kg⁻¹ p.o.) served as reference standard and Group III animals received (0.01 ml Freund's adjuvant) served as an arthritic control and Group IV and V animals received the crude ethanolic extract of *Clerodendrum phlomidis* (250 and 500 mg.kg⁻¹). The ethanolic extract was administered after 14 days from the day of adjuvant injection for 14 days by intubation and the drug treatment was given until 28th day by oral route. All radiographs were taken with a Wipro GE X-ray instrument set at 45kV and 4 mAs using Laser Orthochromic film. The film -to- source distance was 100cm. X-ray was taken at the joints of the hind paw for the confirmation and evaluation of the severity of arthritis in FAI induced rats. The X-ray examination of Hind Limb Bone Mass of animals in HLBM of Group I – Group V was carried out continuously for 5 weeks [17]. On day 1 the animals were subjected to X-ray examination without any treatment and after 3hrs the animals GP I- IV were induced with 0.1ml of Freund's adjuvant in left hind paw of rats. On 7th day X-ray examination of animals in all groups were carried out. On day 14th day X-ray was taken after treatment with ethanolic extract of *Clerodendrum phlomidis* and indomethacin to the corresponding group of animals. The same treatment was continued until 28th day but the X-ray examination was carried out on 21st and 28th day. Estimation of HLBM using photodensitometer and aluminium step wedge. Densitometer is a device by which we can measure the Optical Density (OD) at a particular area of a film and we can measure the density from 0 to 4 with 0.01 accuracy. The processed X-ray film carries the visible image in terms of metallic silver pattern and in other words the degree of blackness is directly related to the amount of silver present. The film darkness directly depend on intensity of radiation reaching the film which inturn depends on atomic number and density of the tissue through which uniform X-ray beam has passed. By measuring the amount of silver loss or degree of blackening, is indicator of nature of tissue. For radiographic standardization x-ray tube voltage KeV, mAs, developing condition should be as a kept constant. A high level of standardization is required

in both projection geometry and image acquisition, which is needed to achieve a precise measurement of density. Radiographic projection geometry is defined by the relative location, and orientation of X-ray source, the object and the film detector. After assessing the nature of tissue, a quantitative measurement was performed using aluminium step wedge. The step of height of aluminium step wedge is 1.5mm to 10.5mm with a width of 3mm was placed on the film cassette and radiograph was exposed. Aluminium was chosen for step wedge since its atomic number is very similar to the effective atomic number of bone. Mineral with similar atomic number will attenuate X-ray in a similar manner. OD of each step of the step wedges was measured and the values were plotted against the corresponding thickness of aluminum. The curve obtained provided the corresponding aluminium equivalent to the measured optical density of the Hind Limb. In this way an indication of the HLBM was obtained.

3. Results

3.1 Acute toxicity

The leaf extract of *Clerodendrum phlomidis* didn't show any mortality and toxicity even at highest dose of 2000 mg.kg⁻¹ body weight employed. The present research study was carried out using two different doses (low and high) ethanolic extract of *Clerodendrum phlomidis* such as 250 and 500mg.kg⁻¹ body weight for anti-arthritis activity.

3.2 PAW VOLUME

An oral administration of *Clerodendrum phlomidis* (250 and 500mg.kg⁻¹) showed a marked inhibition of edema of adjuvant induced chronic arthritic rats and the maximum effect was observed on 28th day and the effect is very similar to the standard drug "Indomethacin". The results are shown in Fig. 2.

3.3 BODY WEIGHT CHANGES

Clerodendrum phlomidis (250 and 500 mg.kg⁻¹) inhibits the loss of body weight on adjuvant induced arthritic animal than compared to vehicle control animals, when the loss of body weight is predominant and indomethacin could ameliorate the weight loss occurred during arthritis. The results are shown in **Table.1**

3.4 X-RAY Analysis (Densitometry)

Radiographs were taken in the left hind paw once in a week and this procedure was followed for 5 weeks before and after treatment of ethanolic extract of *Clerodendrum phlomidis* and standard drug indomethacin. Adjuvant – induced group shows severe bone swelling. The bone density changes were evaluated using photodensitometer and quantitative measurements were made using aluminium step wedge and HLBM was calculated. HLBM of rats was increased by Freund's adjuvant as compared to control group. Interestingly the arthritic animals treated with ethanolic extract of *Clerodendrum phlomidis* (250 and 500 mg.kg⁻¹) showed significant reduction in bone density which is similar to that of standard drug Indomethacin. From the aluminium step wedge we can measure the accurate bone mass density of

control and drug treated groups. There was a statistically significant ($p < 0.001$) difference in mean values of densitometry reading was observed and the value was found to be 0.01. The results of present study indicate that the plant extract treatment successfully suppressed the RA induction. The results are shown in Table.2, Fig.1 and Fig.3.

4. Discussion

Arthritis is a chronic inflammatory disorder and the inflammation involves the release of mediators like cytokines (IL-1B and TNF- α), GM-CSF, interferons and PGDF. These are responsible for the pain, destruction of bone and cartilage that can lead to severe disability [18]. The determination of paw swelling is apparently simple, sensitive and quick procedure for evaluating the degree of inflammation and assessing of therapeutic effects of drugs. In the present study, the rat was selected as an animal model since they develop a chronic swelling in multiple joints with an influence of inflammatory cells and followed by erosion of cartilage in joints and destruction of bones. The rat model is a close resemblance to rheumatoid arthritis of human beings [19]. From our study we observed that the ethanolic extract of *Clerodendrum phlomidis* and indomethacin significantly suppressed the swelling of the paws of rats.

A change in body weight of rats was also measured as one of the parameter to assess the course of the disease and the response to therapy of anti-inflammatory and arthritic drugs [20]. As the incidence and severity of arthritis increased, a decrease in body weights of the rats also occurred during the course of the experimental period and this observation was supported by the findings of C.V.Winder [21] on alterations in the metabolic activities of diseased rats. In addition to the absorption of ¹⁴C- glucose and ¹⁴C- leucine in rat's intestine was reduced in the case of inflamed rats

[22] put on the treatment with anti-arthritic drugs, the decrease in absorption was nullified [23] and it shows that the anti-arthritic drugs correct the decreased/deranged absorption capacity of intestine during arthritis. The increased body weight during treatment of indomethacin and ethanolic extract of *Clerodendrum phlomidis* may be due to the restoration of absorption capacity of intestine. By using image analysis techniques of radiographs, we measured bone swelling, optical density of all groups of tibio tarsal joints of rats using photodensitometer and aluminium step wedge thickness (mm). This method provides a more sensitive and quantitative approach for radiological image analysis as compared with conventional observation. From the study observed a change of bone swelling by measuring OD in terms of aluminium equivalence (mm) after treatment with ethanolic extract of *Clerodendrum phlomidis* and indomethacin treatment. It has been

previously demonstrated that these measurements are positively correlated with the results of conventional radiological and histological evaluation [18]. The radiographic analysis of the tibio tarsal joint in arthritic and drug treated animals further supported and confirms the potent anti-arthritic effect of *Clerodendrum phlomidis* in a dose dependent manner and it suppress the pathological changes such as pannus formation [24] and bone destruction [25].

5. Conclusion

The ethanolic extract of *Clerodendrum phlomidis* has anti-arthritic and peripheral analgesic on acute and possibly chronic inflammatory processes. The claim made by tradipractitioners [26] that *Clerodendrum phlomidis* use to treat various pains and arthritic diseases is found.




Fig.1 Anti-arthritic effect of *Clerodendrum phlomidis* on rats measured using optical densitometer of X-ray films on 28th day where the control animals with normal saline (A); Animal induced with adjuvant and treated with standard drug "Indomethacin" (B) ; Animal induced with arthritis using adjuvant (C); Animal induced with adjuvant and treated with 250mg/kg of *Clerodendrum phlomidis* (D); Animal induced with adjuvant and treated with 500 mg/kg of *Clerodendrum phlomidis* (E); Aluminium foil step wedge to measure the optical bone density (F).




Fig.2. The anti-arthritic effect of ethanolic extract of *Clerodendrum phloimidis* (250 and 500 mg.kg⁻¹ body weight) was tested by Freund's adjuvant paw edema in rat. The indomethacin (10 mg.kg⁻¹ body weight) was used as a standard drug. The control animal was induced with saline (5 ml.kg⁻¹ body weight). The anti-arthritic effect was tested in different time interval such as 1, 4, 8, 12, 16, 20, 24 and 28 days. Each value represents mean \pm S.E.M, n=6. The statistical analysis was carried out using one way ANOVA method, where **P < 0.01, * P < 0.05

Table. 1: Body weight changes in adjuvant induced arthritis in rats

Group	Mean body weight (gm)		Mean changes in body weight (\pm SEM)
	Before induction	After treatment (On 28 th day)	
Control (Normal saline 5 mg.kg ⁻¹)	158.6	166.4	8.33 \pm 1.667
Standard (Indomethacine 10 mg.kg ⁻¹)	155.1	195.5	40 \pm 2.582**
<i>Clerodendrum phloimidis</i> 250 mg.kg ⁻¹	151.2	174.2	23.33 \pm 4.595**
<i>Clerodendrum phloimidis</i> 500 mg.kg ⁻¹	151.8	162.7	10.83 \pm 1.537

Table.1 The anti-arthritic effect of ethanolic extract of *Clerodendrum phloimidis* (250 and 500 mg.kg⁻¹ body weight) was tested by measuring the change of body weight. The Indomethacine (10 mg.kg⁻¹ body weight) was used as a standard drug. The control animal was treated with saline (5ml.kg⁻¹ body weight). The arthritis was induced with 0.1 ml of FCA. The anti-arthritic effect was tested by before induction and after induction of arthritis. Each value represents mean \pm S.E.M, n=6. The statistical analysis was carried out using one way ANOVA method, where **P < 0.001, *P < 0.01.

Treatment	Day – 1		Day - 7		Day - 14		Day - 21		Day – 28	
	O.D	Step wedge Thickness	O.D	Step wedge Thickness	O.D	Step wedge Thickness	O.D	Step wedge Thickness	O.D	Step wedge Thickness
Control	2.8 0±0.0075	1.7± 0.0248	2.9±0.0.236	1.6± 0.00752	2.8±0.00816	1.8± 0.0075	2.9±0.0236	1.9± 0.0136	2.7±0.0075	1.72±0.0248
Standard	2.6±0.0542	1.6± 0.0075	3.6±0.0816	2.16±0.0.00816***	3.4±0.108	2.13±0.0116***	2.8±0.0075	1.89±00089***	2.5±0.0543	1.68±0.01505***
Adjuvant induced Arthritis	2.9±0.0236	1.76±0.0150	3.8±0.0816	2.17±0.00816***	3.7±0.516	2.11± 0.0116*	3.6±0.0547	2.18±0.0081*	3.5±0.0547	2.18±0.00816**
D.Elata 250mg/kg	2.8±0.0075	1.81±0.0075	3.7±0.0516	2.18±0.00812***	3.2±0.109	2.11 ±0.0116*	2.9±0.0236	1.88±0.0075***	2.6±0.0547	1.71 ±0.0248**
D.Elata 500mg/kg	2.8±0.0.00752	1.72±0.0248	3.7±0.00248	2.16± 0.00814***	3.3±0.108	2.14±0.0082***	3±0.1085	1.86 ±0.015**	2.4±0.0150	1.68 ±0.01505***

Table.2 Anti-arthritic effect of ethanolic extract of *Clerodendrum phlomidis* by X-ray image analysis (optical density Vs Step wedge thickness)

The anti-arthritic effect of ethanolic extract of *Clerodendrum phlomidis* (250 and 500 mg.kg⁻¹ body weight) was tested by x-ray image analysis technique in rat. The Indomethacine (10 mg.kg⁻¹ body weight) was used as a standard drug. The control animal was treated with saline (5mg.kg⁻¹ body weight). The arthritis was induced with 0.1 ml of AIA. The anti-arthritic effect was tested in different time interval such as 1, 2, 3, 4 and 5weeks. Each value represents mean ± S.E.M, n=6. The statistical analysis was carried out using one way ANOVA method, where **p< 0.001 **p < 0.01, * p< 0.5




Fig.3.The anti-arthritic effect of ethanolic extract of *Clerodendrum phlomidis* (250 and 500 mg.kg⁻¹ body weight) was tested using X-ray image analysis (optical density) in rat. The Indomethacin (10 mg.kg⁻¹ body weight) was used as a standard drug. The control animal was treated with saline (5 mg.kg⁻¹ body weight). Arthritis was induced with Freund's adjuvant (0.1 ml). The anti-arthritic effect was tested in different time interval such as 1, 2, 3, 4 and 5 weeks. Each value represents mean \pm S.E.M, n=6. The statistical analysis was carried out using one way ANOVA method, where ***P < 0.001, **P<0.01 n=6.

References

1. Firestein G.S. Evolving concepts of rheumatoid arthritis. *Nature (London)* 2003; 423:356–61.
2. Berens D.L, Lockie L.M, Lin R, et al. Roentgen changes in early rheumatoid arthritis. *Radiology* 1964; 82:645 - 653.
3. Koburova K.L, Michailova SG., Shkenderov SV., Further investigation of the anti-inflammatory properties of adolapin-bee venom polypeptide, *Acta Physiol Pharmacol Bulg* 1985;11: 50-5
4. Martin W., Hartter P. Basic peptides in bee venom, VI. Strutr-activity studies on the anti-inflammatory effects of derivatives and fragments of the MCD-peptide. *Hoppe Seylers Z Physical Chem* 1980; 361:525-35.
5. By waters EGL. The early radiological signs of rheumatoid arthritis. *Bull Rheum Dis* 1960; 11:231-234.
6. Feldmann M., Brennan F.M., Maini RN (1996). Rheumatoid arthritis. *Cell* 85: 307-310.
7. Koch A.E (1998). Angiogenesis: implications for rheumatoid arthritis. *Arthritis Rheum.* 41: 951-962.
8. Sarzi-Puttini P., Comi D., Boccassini L., Muzzupappa S., Turiel M., PanniB., Salvaggio A (2000). Diet and rheumatoid arthritis. A controlled double-blind study of two different dietary regimens. *Scand. J. Rheumatol.* 29: 302-307.
9. Kalpan, H.R., Wolke, R.E., Malone. M.H. 1967. Anti-inflammatory evaluation of cryogenin. *Journal of Pharmaceutical Sciences* 56, 1385-1392.
10. Watson, W.C., Malone, M.H., 1977. Evaluation of Ali, B.H., immunosuppressive potential of cryogenine using developing and established adjuvant arthritis in rats. *Journal of Pharmaceutical Sciences* 66, 1304-1308
11. Ali, B.H., Bashir, A., K., Tanira, M.O., 1995. Anti-inflammatory, anti-pyretic and analgesic effects of *Lawsonia inermis* L. (Henna) in rats. *Journal of Pharmacology* 51, 356-363.
12. Ahmad F., Khan R.A., and S. Rasheed. Preliminary screening of methanolic extracts of *Celastrus paniculatus* and *Tecomella undulate* for analgesic and anti-inflammatory activities. *J Ethanopharmacology* 6, 293-301
13. Shan, N.C., 1982. Herbal Folk medicines in Northern India. *J Ethanopharmacology* 6, 293-301
14. Tunon, H., Olavsdotter, C., Bohlin L, 1995. Evaluation of anti- inflammatory activity of some medicinal plants, inhibition of prostaglandin biosynthesis and PAF- induced exocytosis. *J Ethanopharmacology* 48, 61-76.
15. Ecobichon, D.J., 1977. The Basis of Toxicology Testing. CRP Press, New York, pp.43-86.

16. Newbould, B.B., Chemotherapy of arthritis induced in rats by mycobacterial adjuvant. *Br J PharmacolChemother.* 21: 127-36 (1963). Possible mechanism in enhancing cognition .*Phytomedicine.* 9(4): 302-11 (2002).
17. Esser R.E., Hildebrand A.R., Angelo R.A., Watts L.M., Murphey M.D., and Baugh LE. Measurement of radiographic changes in adjuvant-induced arthritis in rats by quantitative image analysis. *Arthritis Rheum* 1995; 38: 129-38 *J Ethnopharmacol.* 193-8 (1994).
18. Eric G.B., Lawrence J.L., Rheumatoid Arthritis and its therapy. The textbook of therapeutics drug and disease management. 16th Ed. (Williamsand Wilkins Company, Baltimore, 1996) pp. 579-95.
19. Singh S., Majumdar D.K. Effect of fixed oil of *Ocimum sanctum* against experimentally induced arthritis and joint edema in laboratory animals .*International Journal of Pharmacognosy* 34(3):218-22 (1996).
20. William J.K., Arthritis and allied condition. A textbook of rheumatology. 3rd Edn Vol.-1. (AWaverlay Company, Baltimore, Tokyo, 1996) pp.1207-26.
21. Winder C.V., Lembke L.A., Stephens M.D. Comparative bioassay of drugs in adjuvant-induced arthritis in rats: flufenamic acid, mefenamic acidand phenylbutazone. *Arthritis Rheum.* 12(5): 472-82 (1969).
22. Walz D.T., Dimartino M.J., and Misher A(1971). Adjuvant-induced arthritis in rats. II. Drug effects on physiologic, biochemical, and immunological parameters. *J Pharmacol Exp Ther.* 178(1): 223-31.
23. Somasundaran S., Sadique J., Subramoniam A. (1983). Influence of extra-intestinal inflammation on the *in vitro* absorption of ¹⁴C-glucose and the effects of anti-inflammatory drugs in the jejunum of rats. *Clin Exp Pharmacol Physiol.* 10(2): 147-52 (1983).
24. Koburova K.L, Michailova SG., Shkenderov SV., Further investigation of the anti- inflammatory properties of adolapin-bee venom polypeptide, *Acta Physiol Pharmacol Bulg* 1985;11: 50-5
25. Simon G. Alteration in joint space (arthritis) and aspirated bone change, in: *principal of Bone X-ray diagnosis*, 2nd ed., Butler worth & co. Ltd., Great Britain, 1965, pp. 157-163.
26. Adjanohoun, J.E., Aboubakar, N., Dramane, K., Ebot, M.E., Ekpere, J.A.,et al., 1996. OAU/STRC Traditional Medicine and Pharmacopoeia: Contribution to Ethnobotanical and Floristic Studies in Cameroon. CNPMS, Porto Novo, Benin, pp. 360–420.
