

ANTICANCER ACTIVITY OF PLANT: *ADIANTUM VENUSTUM* *DON*

¹Devmurari V.P.*, ¹Pandey S., ¹Goyani M.B, ¹Jivani N.P,
²Marotrao S, ²P. Sivakumar

¹Department of Pharmaceutical Chemistry, Smt R B P Mahila Pharmacy College, Atkot,
Ta: Jasadan, Gujarat, India

²Department of Pharmaceutical Chemistry, J.K.K. Nataraja College of Pharmacy,
Komarapalayam, Tamilnadu, India

*Corres.author: viraldev1985@gmail.com
Mob no:09724337397

ABSTRACT: Cancer is a malignant disease that is characterized by rapid and uncontrolled formation of abnormal cells which may mass together to form a growth or tumour, or proliferate throughout the body. Next to heart disease cancer is a major killer of mankind. Present study aims at a preliminary phytochemical screening and anticancer evaluation of *Adiantum venustum* Don against Ehrlich Ascites Carcinoma in animal model. Results indicates that ethanolic extract of *Adiantum venustum* Don possess significant anticancer activity and also reduce elevated level of lipid peroxidation due to higher content of terpenoids and flavonoids. Thus ethanolic extract of *Adiantum venustum* Don could have waste therapeutic application against cancer.

KEY WORDS: Cancer, *Adiantum venustum*, Lipid peroxidation, Flavonoids, Terpenoids

INTRODUCTION

The chemotherapy of neoplastic disease has become increasingly important in recent years. The relatively high toxicity of most anticancer drugs has fostered the development of supplementary drugs that may alleviate this toxic effect or stimulate the regrowth of depleted normal cells. Plants have a long history of use in the treatment of cancer. Plants have played an important role as a source of effective anticancer agent, and it is significant that over 60% of currently used anticancer agents are derived in one way or another from natural sources, including plants, marine organism and microorganisms.¹ It was also observed from Ayurvedic literature and ethanobotanical studies that the plant is very useful in treating tumour,

prevention of hair from falling and as a diuretics but no scientific investigation has been done in such direction.²⁻⁴ Therefore it was thought worth while to carry out preliminary phytochemical screening and screening of *Adiantum venustum* Don for anticancer activity against Ehrlich Ascites Carcinoma in animal model.

EXPERIMENTALS

Plant source

The plant leaves and stem of *Adiantum venustum* Don were collected from Kolli hills, Namakkal District, Tamilnadu, India, and were authenticated. Reference number of the authentication report is BSI/SC/5/23/05.06/Tech/603.



Figure 1: Plant Image (*Adiantum venustum* Don)

Extraction Procedure

The leaves and stem of *Adiantum venustum* were dried under shade, mixed together and then made into a coarse powder with a mechanical grinder. The powder was passed through sieve no. 40 and stored in an airtight container for further use. The dried powder material (150gm) was defatted with petroleum ether (60-80°) to remove waxy substances and chlorophyll, which usually interfere in the isolation of phytoconstituents. The marc after defatted with petroleum ether was dried and extracted with ethanol (99.9 % v/v) in a Soxhlet extractor for 72 hr. The solvent was then distilled off and the resulting semisolid mass was dried in a vacuum evaporator to get a yield of 14 % w/w.^{5,6}

Phytochemical Identification Tests

Various chemical tests were performed for the Phytochemical identification of the ether and ethanolic extract of the plant leaves and stem of *Adiantum venustum* as per standard procedure.^{7,8}

Anticancer activity

Toxicity Evaluation (LD₅₀): (Karber's methods)

Thirty mice including both male and female weighing 20–25 gm were selected for the study. LD₅₀ was measured by Karber's methods

Animals

Male Swiss albino mice weighing between 18-25 gm were used for present study. They were maintained under standard environmental conditions and were fed with standard pellet diet of water and *ad libitum*. The mice were acclimatized and laboratory condition for 10 days before commencement of experiment. All procedure described were reviewed and approved by the Institutional Animal Ethical Committee of J.K.K. Nataraja College of Pharmacy, Komarapalayam.

Cancer Cell line

EAC cells were obtained from Amala Cancer Research Center, Thrissur, Kerala, India. They were maintained by weekly intraperitoneal inoculation of 10⁶ cells / mouse.

Preparation of extract drug and mode of administration

In the present anticancer study, ethanolic extract of *Adiantum venustum* (EEAV) in the dose of 100 mg/kg and 200 mg/kg were prepared as suspension by dissolving the ethanolic extract in propylene glycol and sterile physiological saline containing Tween 20 to get the desired concentration.^{9,10}

Tumor Transplantation

Ehrlich's Ascites Carcinoma was maintained by serial transplantation from tumor bearing Swiss Albino mice. Ascetic fluid was drawn out from tumor bearing mice at the log phase (day 78 of tumor bearing) of the tumor cells. The tumor cell number was adjusted to 2X10⁶ tumor cells/ml. Sample showing more than 90 % viability was used for transplantation. Each animal received 0.2 ml of tumor cell suspension containing 2 X 10⁶ cells / ml intraperitoneally.¹¹

Drug Treatment Schedule:

Male swiss albino mice were divided into 5 groups (n = 8). All the groups were injected with EAC cells (0.2 ml of 2 X 10⁶ cells/mouse) intraperitoneally except the normal group. This was taken as day zero. From the first day normal saline 5 ml/kg/mouse/day and propylene glycol 5 ml/kg /mouse/day was administered to normal and EAC control groups respectively for 14 days intraperitoneally. Similarly EEAV at different doses (100 mg and 200 mg/kg/mouse/day) were administered in groups 3, 4, 5 respectively after the administration of last dose

followed by 18 hrs. fasting 4 mice from each group were sacrificed for the study of antitumor activity, hematological and liver biochemical parameters. The remaining animals in each of the groups were kept to check the mean survival time (MST) and percent increase in life span of the tumor bearing hosts.^{12,13,14} Various parameters like Body weight of animals, Life span of animals, Cytological studies on cell lines, Hematological parameter, RBC, WBC, Hemoglobin, differential count, Biochemical parameters evaluated in the present study.

Anticancer effect of EEAV was assayed by observation of change with respect of body weight, ascitic tumor volume, packed cell volume, viable and non viable tumor cell count, mean survived time (MST) and percentage increase in life span (%ILS).¹²

Tumor Cell Volume and Packed Cell Volume

The mice were dissected to collect ascitic fluid from peritoneal cavity and centrifuged to determine packed cell volume at 1000 rpm for 5 min.¹² The transplantable murrane tumor was carefully collected to measured the tumor volume.

Viable and non viable cell count

Viable and non viable cell counting of the ascetic cell was done by staining with tryphan blue (0.4 % in normal saline), dye exclusion test and count was determined in a Neubauer counting chamber. The cells that did not take up they dye were viable and those that took the stain were not viable.¹²

Mean survival time and percent increased in life span

The effect of EEAV on tumor growth was observed by MST and % ILS. MST of each group continuing 4 mice were monitored by recording the mortality daily for 6 weeks and % ILS was calculated by using following equation.^{11,12}

MST =
(Day of first death + Day of last death)/2

$$\% \text{ ILS} = \left\{ \frac{\text{MST of treated group}}{\text{MST of control group}} \right\} - 1 \times 100$$

Effect of EEAV on hematological parameters

Blood was collected from each mice by intracardial puncture with blood anticoagulant (Heparin) and while blood cells (WBC), red blood cells (RBC); hemoglobin and differential count were determined in group comprise of I) Tumor bearing mice (control), II) Tumor bearing mice treated with EEAV (100 mg/kg/mice/day), III) Tumor bearing mice treated with EEAV (200 mg/kg/mice/day), VI) Normal group.¹⁵

Biochemical Assay

After the collection of blood samples the mice were sacrificed and their liver was excised. The isolated liver was rinsed in ice cold normal saline followed by cold phosphate buffer having pH 7.4, and blotted dry and weighed. A 10% w/v homogenate of liver was prepared in ice cold phosphate buffer (pH 7.4) and a portion were utilized for estimation of lipid peroxidation and other portion of the same after precipitation of proteins with TCA was used for estimation of glutathione remaining homogenate were centrifuged at 1500 rpm at 4°C for 15 min. The supernatant thus obtained was used for the estimation of superoxide dismutase, catalase and protein content.¹⁴

Statistical Analysis

The experimental result were expressed as mean \pm SEM. Data were assessed by the student t-test $P < 0.05$ was considered as statistically significant.

RESULTS AND DISCUSSION

Phytochemical screenings suggest that ethanolic extract of plant contain terpenoid, phytosterols, flavanoid and saponin. (Table 1.)

Anticancer Activity

Toxicity Evaluation (LD₅₀): In acute toxicity study, the given extract of *Adiantum venustum* did not show any mortality up to the dose of 2000 mg / kg. The extract shows sedation, hypnosis, mild muscle relaxant property.

Administration of EEAV reduces the tumour volume, packed cell volume and viable tumour cell count in a dose dependant manner when compared to EAC control mice. In EAC control mice the median survival time was 22 \pm 0.25 days. Whereas, it was significant increased median survival time (24 \pm 0.33, 29 \pm 0.49) with different doses (100 and 200 mg/kg) of EEAV and standard drug respectively. The mean survival time and effect of EEAV (100mg/kg, 200mg/kg) at different doses on tumour volume, viable and non viable cell count, were shown in table 2 & table 3 and graphical representation are shown in following figure.




Figure 2. Graphical representation of mean survival time and % increase in life span of mice.




Figure 3. Graphical representation of effect of EEAV on tumour volume and packed cell volume of mice, viable cell count and non viable cell count




Figure 4. Graphical representation of effect of EEAV Effect on haematological parameters

EAV at the dose of 100 and 200 mg/kg the haemoglobin content in EAC bearing mice were increased to 10.6 ± 0.057 and 11.45 ± 0.057 . The haemoglobin contents in the EAC control mice (9.8 ± 0.02) was significantly decreased as compare to normal mice (12.85 ± 0.25). (table-7) The total WBC count was significantly higher in the EAC treated mice when compared with normal mice. Whereas EAV treated mice significantly reduced the WBC count as compared to that of control mice. Significant changes observed on differential count when extract treated mice compared with EAC control mice. (Table-4)

Biochemical assay

Biochemical assay indicated that EAV significantly reduced the elevated levels of lipid peroxidation and thereby it may act as an antitumor agent. The level of lipid peroxidation, catalase and protein content were summarized in table 8 and graphical representation shown. (Table 5)

The plant leaves and stem of *Adiantum venustum* Don were found to contain higher amount of triterpenoids. In present study anticancer potential of the plant was estimated in EAC bearing carcinoma Cell. Ethanolic extract of *Adiantum venustum* Don had considerably reduced tumour volume and increased the life span of the test animals. It is also found that EAV significantly reduced the elevated levels of lipid peroxidation and thereby it may act as an antitumour agent.

CONCLUSION

The ethanolic extract of *Adiantum venustum* Don possessed significant anticancer and antioxidant activity due to its higher terpenoids and flavonoids content. Further investigation on different biological activities of this plant with different modes will not only validate the types of activities claimed by ayurvedic, siddha and traditional practitioners, but also will bring out innovation in the field of therapeutics.




Figure 5. Graphical representation of effect of EAV on different biochemical parameters

Table 1: Result of phytoconstituent identification tests of ethanol extract of *Adiantum venustum* Don.

Phytoconstituent	Phytosterol	Flavonoids	Triterpenoids	Saponin
Ethanol Extract	+	+	+	+

+ Present - Absent

Table 2 : Effect of Ethanol Extract of *Adiantum venustum* on survival time on EAC bearing mice

S. No.	Experimental groups	Mean survival time (MST) days	% increase in life span
1	Normal control (normal saline 5 ml/kg b.w.)	-	-
2	EAC control	22±0.25	-
3	EAC + EEAV (100 mg /kg)	24±0.33	9.09
4	EAC + EEAV (200 mg / kg)	29±0.49	31.81
5	EAC + Vincristine (0.8 mg / kg) std	31±0.55	40.90

Values are mean ± SEM (Standard error of mean), Number of mice in each group (n=4), P < 0.001, Experimental group was compared with EAC control.

Table 3: Effect of Ethanol extract of *Adiantum venustum* on tumor volume, packed cell volume, viable and non viable tumour cell count of EAC bearing mice.

Parameters	EAC Control	EEAV 100 mg / kg	EEAV 200 mg/kg	Standard vincristine 0.8 mg/kg
Body weight	26.11±0.12	24.34±0.16	23.28±0.13	23.9±0.02
Tumour volume (ml)	5.82±0.042	4.22±0.051	3.42±0.082	2.42±0.13
Packed cell volume (ml)	2.12±0.104	1.75±0.043	1.05±0.092	1.15±0.03
Viable tumour cell count % 10 ⁷ cells /ml	11.25±0.098	7.78±0.18	4.85±0.23	4.90±0.015
Non viable tumour cell count X 10 ⁷ cells / ml	0.5±0.017	0.92±0.023	1.47±0.021	1.23±0.81

Values are mean ± SEM. No. of mice in each group (n = 4), P < 0.01, experimental groups was compared with EAC control Weight of normal mice=20±0.15

Table 4: Effect of Ethanol Extract of *Adiantum venustum* on haematological .parameters of EAC treated mice

Parameter	Normal saline 0.5 ml/kg	EAC control 2 X 10 ⁶ cells / mice	EAC + EEAV 100 mg/kg	EAC + EEAV 200 mg / kg	EAC Cell + Vincristine 0.8 mg/kg
Haemoglobin (gm)	12.85±0.25	9.8±0.02	10.6±0.057	11.45±0.18*	11.7±0.045*
Total RBC million/mmcu	6.65±0.18	3.8±0.035	4.75±0.032	5.42±0.22*	5.8±0.054
Total WBC Million/mmcu	7.8±0.045	20.07±0.068*	11.92±0.042	8.85±0.059	9.12±0.055
Lymphocyte	77.75±0.19	33.37±0.56*	52.7±0.50*	60.72±0.36*	59.12±0.30
Monocyte	1.7±0.035	0.82±0.024	1.15±0.014*	1.2±0.045	1.32±0.024
Granulocyte	29.97±0.46	52.6±0.37*	40.87±0.2	31.72±0.63*	41.65±0.29

Values are mean ±SEM , (n=4), EAC control group compared with normal group, Experimental group compared with EAC control. P < 0.01, *P < 0.05

Table 5 : Effect of different doses of ethanolic extract of *Adiantum venustum* on different biochemical parameter in EAC bearing mice.

Parameter	Normal saline 0.5 ml/kg	EAC control 2 X 10 ⁶ cells / mice	EAC + EEAV 100 mg/kg	EAC + EEAV 200 mg / kg
Lipid peroxidation n mole MDA/gm of tissue	0.92±0.02	1.36±0.09*	1.27±0.04*	1.13±0.02
Catalase (units /mg tissues)	2.51±0.72	1.71±0.15*	1.75±0.13	2.34±0.23*
Protein content (gm / 100 ml)	12.66±0.69*	17.25±0.76	16.50±0.70	16.10±0.55

Values are mean ±SEM, (n=4)EAC control group compared with normal group, Experimental group compared with EAC control. P < 0.05, *P< 0.05

REFERENCES

- Kadam S.S., Mahadik K.R., Bothra K.G, "Principle of Medicinal Chemistry", 1st edition, Nirali Prakashan, 1989, 86.
- Kirtikar KP, Basu BD, "Indian Medicinal plants", IInd ed, Vol IV, 1935,2738-2739.
- Natkarni's K.M., "Indian Materia Medica", vol I,1976, 44.
- Ambarta SP, The Useful plants of India, CSIR, New Delhi, 1986, 15.
- Harbone JB, "Phytochemical Method, A Guide to modern techniques of plant Analysis", 3rd edition, springer (Indian) pvt. ltd., New Delhi, 2005, 16,-22.
- Krishnaswamy NR. Chemistry of Natural products, A laboratory hand book, 1st edition, Universities press India (pvt.) Ltd, Hyderabad, 2003, 15, 26-30, 70-73, 87-88.
- Mohammad A, Objective type pharmacy, 2nd edi, Birla publication, Delhi, 2000-2001, 567-568.
- Khandelwal KR, Kokate CK, Pawar AP, Gokhle SB, Practical pharmacognosy Techniques and Experiments, 3rd edition, Nirali prakahan, Pune, 1996,165-166.
- Teresa K, Joseph S, Preparative Layer Chromatography, chromatographic science series, 1995.
- Ng TB, Gao W, Li L, Niu SM,Zhao L, Liu J, Shi LS, Fu M , Liu F, Rose (*Rosa rugosa*) – flower extract increases the activities of antioxidant enzymes and their gene expression and reduce lipid peroxidation. Biochem. Cell Biol., 2005; 83: 78-85
- Nicol BM, Prasad SB, The effect of cyclophosphamide alone and in combination with ascorbic acid against murine ascites Dalton's lymphoma. Indian J. Pharmacol., 2006, 38(4), 260-265.
- Sivakumar T, Sambathkumar R, Perumal P, Vamsi MLM, Sivakumar , Kanagasabai R, BaskaranMV, Subhas SK, Mazumdar UK, and Gupta M, Antitumor and antioxidant activity of *Bryonia laciniosa* against Ehrlich's Ascites Carcinoma bearing Swiss Albino mice., Oriental Phar. and Exp. Med., 2005, 5(4), 322-330.
- Kavitha K, Manoharan S, Anticarcinogenic and antilipidperoxidative effect of *Tephrosia purpurea* (Linn). Pers, in 7,12- dimethyl benz (a) anthracene (DMBA) induced hamster buccal pouch carcinoma., Indian J. Pharmacol., 2006, 38(3), 185-189.
- Gupta M, Mazumdar UK, Sambathkumar R, Sivakumar T, "Antitumor activity and antioxidant role of *Bauhinia racemosa* against Ehrlich ascites carcinoma in swiss albino mice", Acta Pharm. Sinica., 2004,25(8), 1070-1076.
- Khanam JA, Bag SP, Sur B, Sur P, Antineoplastic activity of copper benzohydroxamic and complex against Ehrlich ascites carcinoma (EAC) in mice. Indian J. Pharm., 1997, 29(3),157-161.
