

Hepatoprotective activity of some indigenous plants

Sharma Bhawna* , Sharma Upendra Kumar

Institute of pharmacy, Bundelkhand University, Kanpur road,

Jhansi, India.

*Corres.author: bms1928@yahoo.com
Mobile no 9793044783, 9935655434

Abstract: Herbal plants or botanical medicines have been used traditionally by herbalist worldwide for the prevention and treatment of liver disease. The plant kingdom plays a major role in the life of human beings and animals. The plant, as one of the important sources, still maintains its original place in the treatment of various diseases, including liver disorders, with no ill effects. Considerable studies have been carried out on ethno medicinal plants; however, only few medicinal plants have attracted the interest of scientists, to investigate them for a remedy for hepatoprotective. Clinical research in this century has confirmed the efficacy of several plants in the treatment of liver disease. Hence, this review article contributes to the knowledge of reported indigenous plants, which are prevalent for prevention and treatment of liver disorders.

Key words: Hepatoprotective plants, liver, Pharmacognosy, Pharmacological activities, Natural products, Medicinal herbs.

Introduction

The liver is the largest glandular organ in the body, and has more functions than any other human organ. A person's entire blood supply passes through the liver several times a day; The Liver has a pivotal role in human metabolism. Liver produces and secretes bile, it also produces prothrombin and fibrinogen, both blood-clotting factors, and heparin, a mucopolysaccharide sulfuric acid ester that helps keep blood from clotting within the circulatory system. The liver converts sugar into glycogen,

Liver diseases have become one of the major causes of morbidity and mortality in man and animals all over globe and hepatotoxicity due to drugs appears to be the most common contributing factor¹. Among the many diseases that can affect the liver the most common is 'viral hepatitis' (Inflammation of liver caused by viral infection). Hepatitis can be caused by drugs, viruses, bacteria, mushrooms, parasites like amoebas or giardiasis. About 20,000 deaths found every year due to liver disorders. The use of natural remedies for the treatment of liver diseases has a long history and medicinal plants and their derivatives are still used all over the world in one form or the other for this purpose. Scientific evaluation of plants has often shown that active principles in these are responsible for therapeutic success. A large number of medicinal plants have been tested and found to contain active

principles with curative properties against a variety of diseases². Liver protective plants contain a variety of chemical constituents like phenols, coumarins, lignans, essential oil, monoterpenes, carotinoids, glycosides, flavanoids, organic acids, lipids, alkaloids and xanthenes³. Recent experience has shown that plant drugs are relatively non-toxic, safe and even free from serious side effects⁴. This review article has been presented to enumerate some indigenous plants that have hepatoprotective properties.

Hepatoprotective plants-

***Andrographis lineata* nees** Hepatoprotective effect of *Andrographis lineata* (Acanthaceae) extracts in CCl(4)- induced liver injury in rats. Male Wistar rats with chronic liver damage, induced by subcutaneous injection of 50% v/v CCl(4) in liquid paraffin at a dose of 3 mL/kg on alternate days for a period of 4 weeks, were treated with methanol and aqueous extracts of *A. lineata* orally at a dose of 845 mg/kg/day. The biochemical parameters such as serum glutamate oxaloacetate transaminase, serum glutamate pyruvate transaminase, serum bilirubin and alkaline phosphatase were estimated to assess the liver function. Histopathological examinations of liver tissue corroborated well with the biochemical changes. The

activities of extracts were comparable to a standard drug⁵.

Andrographis paniculata (kalmegh) Antihepatotoxic activity of the *Andrographis paniculata* (acanthaceae) methanolic extract (equivalent to 100 mg/kg of andrographolide) and 761.33 mg/kg ip, of the andrographolide-free methanolic extract (equivalent to 861.33 mg/kg of the methanolic extract) of the plant, using CCl₄-intoxicated rats. Biochemical parameters like serum transaminases--GOT and GPT, serum alkaline phosphatase, serum bilirubin and hepatic triglycerides were estimated to assess the liver function. The results suggest that andrographolide is the major active antihepatotoxic principle present in *A. paniculata*⁶.

Azadirachta indica (Neem) Effect of *A. indica* leaf (*meliaceae*) extract on serum enzyme levels (glutamate oxaloacetate transaminase, glutamate pyruvate transaminase, acid phosphatase and alkaline phosphatase) elevated by paracetamol in rats was studied with a view to observe any possible hepatoprotective effect of this plant. It is stipulated that the extract treated group was protected from hepatic cell damage caused by paracetamol induction. The findings were further confirmed by histopathological study of liver. The antihepatotoxic action of picroliv seems likely due to an alteration in the biotransformation of the toxic substances resulting in decreased formation of reactive metabolites⁷.

Careya arborea The methanol extract of *Careya arborea* bark, (*myrtaceae*) was tested for antioxidant and hepatoprotective activity in Ehrlich ascites carcinoma (EAC) tumor-bearing mice. Tumor control animals inoculated with EAC showed a significant alteration in the levels of antioxidant and hepatoprotective parameters⁸. The extract treatment at 50, 100 and 200 mg/kg body weight doses given orally caused a significant reversal of these biochemical changes towards the normal in serum. Liver and kidney when compared to tumor control animals indicating the potent antioxidant and hepatoprotective nature of the standardized extract.

Cassia fistula (Amaltas) Hepatoprotective activity of the n-heptane extract of *Cassia fistula* (*Fabaceae*) leaves was investigated by inducing hepatotoxicity with paracetamol in rats. The extract at a dose of 400 mg/kg body wt. exhibited orally, significant protective effect by lowering the serum levels of transaminases (SGOT and SGPT), bilirubin and alkaline phosphatase (ALP). The effects produced were comparable to that of a standard hepatoprotective agent⁹.

Cleome viscosa Linn (Tickweed) The hepatoprotective activity of the *Cleome viscosa* Linn (*Capparidaceae*) extract was assessed in CCl₄ induced hepatotoxic rats.

The test material was found effective as hepatoprotective, through in vivo and histopathological studies. The extract was found to be effective in shortening the thiopental induced sleep in mice poisoned with CCl₄. The hepatoprotective effect of ethanolic extract was comparable to that of silymarin, a standard hepatoprotective agent¹⁰.

Eclipta Alba (Bhringaraj) The hepatoprotective effect of the ethanol/water (1:1) extract of *Eclipta Alba* (*Asteraceae*) was studied at subcellular levels in rats against (CCl₄) -induced hepatotoxicity. The loss of hepatic lysosomal acid phosphatase and alkaline phosphatase by (CCl₄) was significantly restored by *Eclipta Alba*. The study shows that hepatoprotective activity of *Eclipta Alba* is by regulating the levels of hepatic microsomal drug metabolising enzymes¹¹.

Fumaria indica (hauskn) *Fumaria indica* (*Fumariceae*) were studied for their hepatoprotective activity against carbontetrachloride, paracetamol and rifampicin-induced hepatotoxicities in albino rats. The petroleum ether extract against carbonetrachloride, total aqueous extract against paracetamol and methanolic extract against rifampicin-induced hepatotoxicities showed similar reductions in the elevated levels of some of the serum biochemical parameters in a manner similar that of silymarin indicating its potential as a hepatoprotective agent¹².

***Morinda citrifolia* L. (noni)** The hepatoprotective effects of Noni juice (TNJ) (*Rubiaceae*) against CCl₄-induced chronic liver damage in female Sprague Dawley (SD) rats. Histopathological examination revealed that liver sections from the TNJ + CCl₄ appeared similar to controls, whereas typical hepatic steatosis was observed in the placebo + CCl₄ group. Serum alkaline phosphatase (ALP), aspartate aminotransferase (AST), alanine transaminase (ALT), total cholesterol (TC), triglycerides (TG), low-density lipoprotein (LDL), and very low-density lipoprotein (VLDL) levels were increased in the placebo group compared with the TNJ group. In contrast, high-density lipoprotein (HDL) was increased in the TNJ group and decreased in the placebo group. Thus, TNJ juice appears to protect the liver from chronic exogenous CCl₄ exposures¹³.

Phyllanthus amarus (Bhuiamala) Ethanolic extract of *Phyllanthus amarus* (*euphorbiaceae*), at (0.3g kg (-1) BW 0.2 ml (-1) day (-1) was given to all groups except control groups (gp. I and gp. V), after 30 min of aflatoxin administration. The entire study was carried out for 3 months and animals were sacrificed after an interval of 30 days till the completion of study. *Phyllanthus amarus* extract was found to show hepatoprotective effect by lowering down the content of thiobarbituric acid reactive substances (TBARS) and enhancing the reduced glutathione level and the

activities of antioxidant enzymes, glutathione peroxidase (GPx), glutathione-S-transferase (GST), superoxide dismutase (SOD) and catalase (CAT)¹⁴.

Phyllanthus emblica (amla) Ethanol extract of *Phyllanthus emblica* Linn. (Euphorbiaceae) (PE) induced rat hepatic injury. PE (0.5 and 1 mg/ml) increased cell viability of rat primary cultured hepatocytes being treated with ethanol (96 μ l/m) by increasing % MTT and decreasing the release of transaminase. Pretreatment of rats with PE at oral dose of 25, 50 and 75 mg/kg or SL (silymarin, a reference hepatoprotective agent) at 5 mg/kg, 4 h before ethanol lowered the ethanol induced levels of AST, ALT and IL-1 β . The 75 mg/kg PE dose gave the best result similar to SL. Treatment of rats with PE (75 mg/kg/day) or SL (5 mg/kg/day) for 7 days after 21 days with ethanol (4 g/kg/day, p.o.) enhanced liver cell recovery by bringing the levels of AST, ALT, IL-1 β back to normal¹⁵.

Phyllanthus polyphyllus Methanolic extract of *Phyllanthus-polyphyllus* (euphorbiaceae) was evaluated for hepatoprotective and antioxidant activities in rats. The plant extract (200 and 300 mg/kg, p.o.) showed a remarkable hepatoprotective and antioxidant activity against acetaminophen induced hepatotoxicity as judged from the serum marker enzymes and antioxidant levels in liver tissues¹⁶. The activity of the extract at dose of 300 mg/kg was comparable to the standard drug, silymarin (50 mg/kg, p.o.). Histopathological changes of liver sample were compared with respective control.

Phyllanthus reticulatus (Potato bush) Two partially purified organic fractions designated by PR1 and PR2 of the fat free ethanol (95%) extract of aerial parts of *Phyllanthus reticulatus* (Euphorbiaceae), were tested for the hepatoprotective activity in rats against CCl₄-induced liver damage. The rats receiving the fractions showed promising hepatoprotective activity as evident from significant changes of pentobarbital-induced sleeping time, changes in serum levels of sGPT, sGOT, sALP and bilirubin and also from histopathological changes as compared to CCl₄-intoxicated rats¹⁷.

Picrorhiza kurroa (Kutki) Administration of picroliv, a standardized fraction of alcoholic extract of *Picrorhiza kurroa* (Scrophulariaceae) (3-12 mg/kg/day for two weeks) simultaneously with P. berghei infection showed significant protection against hepatic damage in *Mastomys natalensis*. The increased levels of serum glutamate oxaloacetate transaminase (GOT), glutamate pyruvate transaminase (GPT), alkaline phosphatase, lipoprotein-X (LP-X) and bilirubin in the infected animals were marked reduced by different doses of picroliv. In the liver, picroliv decreased the

levels of lipid peroxides and hydroperoxides and facilitated the recovery of superoxide dismutase and glycogen^{18, 19}.

Polygala arvensis Chloroform extract of leaves of *Polygala arvensis* (polygalaceae), in 0.3% carboxy methyl cellulose (CMC) was evaluated for hepatoprotective activity in Wistar albino rats by inducing hepatic injury with d-galactosamine (400 mg/kg). The chloroform extract of *Polygala arvensis* at an oral dose of 200 mg/kg and 400 mg/kg exhibited a significant ($P < 0.001$, $P < 0.01$ and $P < 0.05$) protection effect by normalizing the levels of aspartate amino transferase (ASAT, GOT), alanine amino transferase (ALAT, GPT), alkaline phosphatase (ALP), total bilirubin (TB), lactate dehydrogenase (LDH), total cholesterol (TC), triglycerides (TGL), albumin, total protein (TP) which were significantly ($P < 0.001$) increased in rats by treatment with 400 mg/kg i.p. of d-galactosamine. Silymarin (25 mg/kg), a known hepatoprotective drug used for comparison exhibited significant activity ($P < 0.001$)²⁰.

Pterocarpus santalinus (Kanak champa) the aqueous (45 mg/ml) and ethanol (30 mg/ml) extracts of *Pterocarpus santalinus* (Fabaceae) stem bark in 1% gum tragacanth was administered orally for 14 days and the hepatoprotective activity studied in CCl₄ induced hepatic damage model. There was a significant increase in serum levels of bilirubin, alanine transaminase, aspartate transaminase and alkaline phosphatase with a decrease in total protein level, in the CCl₄ treated animals, reflecting liver injury. In the extracts treated animals there was a decrease in serum levels of the markers and significant increase in total protein, indicating the recovery of hepatic cells. Ethanol extract treated animal's revealed normal hepatic cords without any cellular necrosis and fatty infiltration²¹.

Pterospermum acerifolium (Kanak champa) the hepatoprotective activity of the ethanol extract of the leaf of *Pterospermum acerifolium* (Sterculiaceae) was investigated in rats. Hepatotoxicity was induced in male Wistar rats by intraperitoneal injection of carbon tetrachloride (0.1 ml/kg/d p.o. for 14 d). Ethanol extract of *P. acerifolium* leaves were administered to the experimental rats (25 mg/kg/d p.o. for 14d). The Hepatoprotective effect of these extracts was evaluated by liver function biochemical parameters (total bilirubin, serum protein, alanine aminotransaminase, aspartate aminotransaminase and alkaline phosphates activities) and histopathological studies of liver. In ethanol extract-treated animals, the toxicity effect of carbon tetrachloride was controlled significantly by restoration of the levels of serum bilirubin and enzymes as compared to the normal and standard drug silymarin-treated groups²².

Solanum nigrum (Makoi) and ***Cichorium intybus*** (Kasni) The presence of plant extracts of *Solanum nigrum* (solanaceae) and *Cichorium intybus* (Asteraceae) in the reaction mixture containing calf thymus DNA and free radical generating system protect DNA against oxidative damage to its deoxyribose sugar moiety. The effect was dependent on the concentration of plant extracts. However, the effect of *Cichorium intybus* was much pronounced as compared to the effect of *Solanum nigrum*. These studies suggest that the observed hepatoprotective effect of these crude plant extracts may be due to their ability to suppress the oxidative degradation of DNA in the tissue debris²³.

Swertia Chirata (Chirayata) Simultaneous treatments with *S. Chirata* (Gentianaceae). (in different doses, viz, 20, 50, and 100 mg/kg body wt daily) and (CCl₄)

caused improvement at both biochemical and histopathological parameters compared to that of (CCl₄) treatment alone but it was most effective when *S. chirata* was administered in a moderate dose (50 mg/kg body wt)^{24, 25}.

Wedelia calendulacea L (Bhanra) Hepatoprotective activity of the ethanolic-leaf extract of *W. calendulacea* (Asteraceae) (EEWC) was studied by estimating serum enzyme activities of aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), protein and bilirubin. 4 The treatment with EEWC showed a dose-dependent reduction of CCl induced elevated serum levels of enzyme activities with parallel increase in total protein and bilirubin, indicating the extract could preserve the normal functional status of the liver²⁶.

References

1. Nadeem M.P.C, Dandiya P.C, K.V., Pasha M., Imran D., Balani K, Vohora S.B., Hepatoprotective activity of *Solanum nigrum* fruits. *Fitoterapia.*, 1997, 68, 245:51.
2. Lewis H.W& Elvin-Lewis M.P.H. Medical Botany: Plants Affecting Man's Health. John Wiley and Sons, New York. 1977, 217-18.
3. Sharma S.K., Ali M and Gupta J., plants having Hepatoprotective activity. *Phytochemistry and Pharmacology.*, 2002, 2, 253-70.
4. Momin A., (1987): Role of indigenous medicine in primary health care. 1st International Seminar on Unani Medicine, New Delhi, India. 1987, 54.
5. Sangameswaran B, Reddy T.C. Jayakar B., (2008): Hepatoprotective effect of leaf extracts of *Andrographis lineata* nees on liver damage caused by carbon tetrachloride in rats. *Phytother Res.*, 2008, 22(1), 124-6.
6. Handa S.S and Sharma A., (1990): Hepatoprotective activity of andrographolide from *Andrographis paniculata* against carbontetrachloride. *Indian J Med Res.*, 1990, 92, 276-83.
7. Chattopadhyay R.R, Sarkar S.K, Ganguly S, Banerjee R.N, Basu T.K, Mukherjee A., Hepatoprotective activity of *Azadirachta indica* leaves on paracetamol induced hepatic damage in rats. *Indian J Exp Biol.*, 1992, 30(8), 738-40.
8. Senthilkumar N, Badami S, Dongre S.H, Bhojraj S., Antioxidant and hepatoprotective activity of the methanol extract of *Careya arborea* bark in Ehrlich ascites carcinoma-bearing mice. *J. Ethnopharmacol.*, 2008, 116(1), 1-6.
9. Bhakta T, Banerjee S, Subhash C. M, Tapan K. Maity, Saha B.P, Pal M., Hepatoprotective activity of *Cassia fistula* leaf extract. *Phytomedicine.*, 2001, 8(3), 220-24.
10. Gupta N.K, Dixit V.K., Evaluation of hepatoprotective activity of *Cleome viscosa* Linn. Extract. *Indian journal of pharmacology.*, 2009, 41, 36-40.
11. Saxena A.K, Singh B, Anand K.K., Hepatoprotective effect of the ethanol/water extract of *Eclipta Alba*. *J Ethnopharmacol.*, 1993, 40(3), 155-61.
12. Rao K.S, Mishra S.H., Hepatoprotective activity of the whole plants of *Fumaria indica*. *Indian journal of pharmaceutical science.* 1997, 59(4), 165-70.
13. Wang M.Y, Anderson G, Nowicki D, Jensen J., Hepatic protection by noni fruit juice against CCl₄-induced chronic liver damage in female SD rats. *Plant Foods Hum Nutr.* 2008, 63(3), 141-5.
14. Naaz F, Javed S, Abidin M.Z., Hepatoprotective effect of ethanolic extract of *Phyllanthus amarus* Schum. et Thonn. On aflatoxin B1-induced liver damage in mice. *J. Ethnopharmacol.*, 2007, 113(3), 503-9.
15. Pramyothin P, Samosorn P, Pongshompoo S and Chaichantipyuth C., the protective effects of *Phyllanthus emblica* Linn. Extract on ethanol induced rat hepatic injury. *J Ethnopharmacol.*, 2006, 107(3), 361-64.
16. B.R, Y.V, J.A, N.H, M.G, V.R., Protective effect of *Phyllanthus polyphyllus* on acetaminophen induced hepatotoxicity in rats. *Pakistan journal of pharmaceutical science.*, 2008, 21(1), 57-62.

17. Das B.K, Bepary S, Datta B.K, Chowdhury A.A, Ali M.S, Rouf A.S., Hepatoprotective activity of *Phyllanthus reticulatus*. Pakistan Journal of Pharmaceutical Sciences., 2008, 21(4), 333-7.
18. Chander R, Dwivedi Y, Rastogi R, Sharma S.K, Garg N.K, Kapoor N.K, Dhawan B.N., Evaluation of hepatoprotective activity of picroliv from *Picrorhiza kurroa* in *Mastomys natalensis* infected with *Plasmodium berghei*. I J Medicinal Res., 1990, 92, 34-7.
19. Ansari R.A, Tripathi S.C, Patnaik G.K, Dhawan B.N., Antihepatotoxic properties of picroliv: an active fraction from rhizomes of *Picrorhiza kurroa*. J Ethnopharmacol., 1991, 34(1), 61-8.
20. Dhanabal S.P, Syamala G, Satish Kumar M.N, Suresh B., Hepatoprotective activity of the Indian medicinal plant *Polygala arvensis* on D-galactosamine-induced hepatic injury in rats. Fitoterapia., 2006, 77(6), 472-4.
21. Manjunatha B.K., Hepatoprotective activity of *Pterocarpus santalinus* L.f., an endangered medicinal plant. Indian journal of pharmacology., 2006, 38, 25-28.
22. Kharpate S, Vadnerkar G, Jain D, Jain S., hepatoprotective activity of the ethanol extract of the leaf of *Portulaca oleraceae*. Indian journal of pharmaceutical sciences., 2007, 69(6), 850-52.
23. Sultana S, Perwaiz S, Iqbal M, Athar M., Crude extracts of hepatoprotective plants, *Solanum nigrum* and *Cichorium intybus* inhibit free radical-mediated DNA damage. Journal of Ethnopharmacol., 1995, 45(3), 189-92.
24. Karan M, Vasisht K, Handa S.S., Antihepatotoxic activity of *Swertia chirata* on paracetamol and galactosamine induced hepatotoxicity in rats. Phytotherapy Research., 1999, 13(2), 95-01.
25. Mukherjee S, Sur A, Maiti B.R., Hepatoprotective effect of *Swertia chirata* on rat. Indian J Exp Biol., 1997, 35(4), 384-8.
26. Murugaian P, Ramamurthy V, and Karmegam N., Hepatoprotective Activity of *Wedelia calendulacea* L. Against Acute Hepatotoxicity in Rats. Research Journal of Agriculture and Biological Sciences., 2008, 4(6), 685-87.
